

Thermochemistry of ammonium based ionic liquids: Thiocyanates - Experiments and computations

Yermalayeu A., Zaitsau D., Emel'yanenko V., Verevkin S.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© Springer Science+Business Media New York 2015. Abstract Molar enthalpies of solution of tetra-n-butylammonium thiocyanate $[N(Bu)_4][SCN]$ and tetra-n-pentylammonium thiocyanate $[N(Pe)_4][SCN]$ in water were measured by using solution calorimetry. The enthalpy of combustion of $[N(Bu)_4][SCN]$ was measured by using rotation bomb combustion calorimetry and the enthalpy of formation of this ionic liquids was derived. The thermal behavior of $[N(Bu)_4][SCN]$ was studied using differential scanning calorimetry. Quantum-chemical calculations of the molar enthalpy of formation in the gaseous phase have been performed for the series $[N(R)_4][SCN]$ with $R = (Me, Et, n-Bu, \text{ and } n-Pe)$ using the G3MP2 level of theory. Experimental and calculated values of the enthalpies of formation are in agreement within the boundaries of the experimental uncertainties.

<http://dx.doi.org/10.1007/s10953-015-0316-2>

Keywords

Combustion calorimetry, Enthalpy of formation, Enthalpy of solution, Ionic liquids, Quantum-chemical calculations, Solution calorimetry